

**Table 101.1 PERIODICITY SCHEDULE FOR  
HEALTH ASSESSMENT REQUIREMENTS BY AGE GROUPS**

Screening Requirement	Age of Person Being Screened														
	Under 1 mo.	1-2 mos.	3-4 mos.	5-6 mos.	7-9 mos.	10-12 mos.	13-15 mos.	16-23 mos.	2 Yr.	3 Yr.	4-5 Yr.	6-8 Yr.	9-12 Yr.	13-16 Yr.	17-20 Yr.
<b>Interval Until Next Exam</b>	1 mo.	2 mo.	2 mo.	2 mo.	3 mos.	3 mos.	3 mos.	6 mos.	1 yr.	1 yr.	2 yr.	3 yr.	4 yr.	4 yr.	None
<b>History and Physical Examination</b>	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
Dental Assessment	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
Nutritional Assessment	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
Developmental/Behavioral	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
Anticipatory Guidance	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
Tobacco Assessment	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
Pelvic Exam 1															•
<b>Measurements</b>															
Head Circumference	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
Height/Length and Weight	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
Blood Pressure											•	•	•	•	•
<b>Sensory Screening</b>															
Visual Acuity Test (Snellen) 2										•	•	•	•	•	•
Clinical Observation	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
Audiometric 2										•	•	•	•	•	•
Non-audiometric	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
<b>Procedures/Tests</b>															
Tuberculin Test											•	•	•	•	•
TB Exposure Risk Assessment	•	•	•	•	•	•	•	•	•	•	•	•	•	•	•
Hematocrit or Hemoglobin					•		•		•	•	•	•	•	•	•
Urine Dipstick or Urinalysis											•	•	•	•	•
Blood Lead Test						•			•						
Blood Lead Risk Assessment					•	•	•	•	•	•	•	•			
<b>Other Laboratory TESTS</b>															
VDRL, RPR, or ART															
Gonorrhea Test															
Chlamydia Test															
Papanicolaou (Pap) Smear															
Sickle Cell															
Ova and Parasites															
<b>Immunizations</b>	Administer as necessary to make status current.														

**NOTE:** Children coming under care who have not received all the recommended procedures for an earlier age should be brought up-to-date as appropriate.

1. Recommended for sexually active females and females age 18 years and older.
2. Snellen testing and audiometric testing should start at age 3 years if possible. Clinical observation and nonaudiometric testing may be substituted if child is uncooperative.